



## Ivermax® (ivermectin) Equine Paste

### Dewormer

#### Product Description

An effective, broad-spectrum paste that helps to control and kill internal parasites, including bots and the arterial stages of *Strongylus vulgaris*, with a single dose. Ivermax Equine Paste contains ivermectin 1.87% anthelmintic and boticide. Apple flavored for palatability.

#### Dosage and Administration

This syringe contains sufficient paste to treat one 1250-pound horse at the recommended dose rate of 91 mcg ivermectin per pound (200 mcg/kg) body weight. Each weight marking on the syringe plunger delivers enough paste to treat 250 pounds of body weight. While holding plunger, turn the knurled ring on the plunger 1/4 turn to the left and slide it so the side nearest the barrel is at the prescribed weight marking. Lock the ring in place by making a 1/4 turn to the right. Make sure that the horse's mouth contains no feed. Remove the cover from the tip of the syringe. Insert the syringe tip into the horse's mouth at the space between the teeth. Depress the plunger as far as it will go, depositing paste on the back of the tongue. Immediately raise the horse's head for a few seconds after dosing.

#### Storage

Store between 20°C - 25°C (68°F - 77°F).


Excursions between 15°C - 30°C (59°F - 86°F) are permitted.

#### Species



Horses

MPN	SIZE	UNIT MEASURE	CASE
21317951	6.08 g	Box of 12 tubes	6 boxes
21302906	6.08 g	Individual tube	100 tubes



See second page for complete indications, precautions, warnings, and residue information.

26FC2001801 (02/25)

Page 1 of 2

# Product Detailer

**Warning:** Do not use in horses intended for human consumption.

Not for use in humans. Keep this and all drugs out of reach of children. Refrain from smoking and eating when handling. Wash hands after use. Avoid contact with eyes. The Material Safety Data Sheet (MSDS) contains more detailed occupational safety information. To report adverse reactions in users, to obtain more information or to obtain MSDS, contact Aspen at 1-816-415-4324.

**Precautions:** Ivermax® (ivermectin) Equine Paste Dewormer has been formulated specifically for use in horses only. This product should not be used in other animal species as severe adverse reactions, including fatalities in dogs, may result.

**Environmental Safety:** Ivermectin and excreted ivermectin residues may adversely affect aquatic organisms. Do not contaminate ground or surface water. Dispose of the syringe in an approved landfill or by incineration.

**Information for Horse Owners:** Swelling and itching reactions after treatment with Ivermax® (ivermectin) Equine Paste Dewormer have occurred in horses carrying heavy infections of neck threadworm (*Onchocerca* sp. *microfilariae*). These reactions were most likely the result of microfilariae dying in large numbers. Symptomatic treatment may be advisable.

Consult your veterinarian should any such reactions occur. Healing of summer sores involving extensive tissue changes may require other appropriate therapy in conjunction with treatment of Ivermax® (ivermectin) Equine Paste Dewormer. Reinfection, and measures for its prevention, should also be considered. Consult your veterinarian if the condition does not improve.

**Indications:** Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism. Ivermax® (ivermectin) Equine Paste Dewormer provides effective treatment and control of the following parasites in horses. *Large Strongyles* (adults) – *Strongylus vulgaris* (also early forms in blood vessels), *S. edentatus* (also tissue stages), *S. equinus*, *Triodontophorus* spp. including *T. brevicauda* and *T. serratus* and *Craterostomum acuticaudatum*; *Small Strongyles* (adults, including those resistant to some benzimidazole class compounds) – *Coronocylus* spp. including *C. coronatus*, *C. labiatus* and *C. labratus*, *Cyathostomum* spp. including *C. catinatum* and *C. pateratum*; *Cylicocylus* spp. including *C. insigne*, *C. leptostomum*, *C. nassatus*, and *C. brevicapsulatus*; *Cylicodontophorus* spp.; *Cylicostephanus* spp. including *C. calicatus*, *C. goldi*, *C. longibursatus*, and *C. minutus* and *Petrovinema poculatum*; *Small Strongyles* – Fourth-stage larvae; Pinworms (adults and fourth-stage larvae) – *Oxyuris equi*; Ascarids (adults, third-stage, and fourth-stage larvae) – *Parascaris equorum*; Hairworms (adults) – *Trichostrongylus axei*; Large-mouth Stomach Worms (adults) – *Habronema muscae*; Bots (oral and gastric stages) – *Gasterophilus* spp. including *G. intestinalis* and *G. nasalis*; Lungworms (adults and fourth-stage larvae) – *Dictyocaulus arnfieldi*; Intestinal Threadworms (adults) – *Strongyloides westeri*; Summer Sores caused by *Habronema* and *Draschia* spp. cutaneous third-stage larvae; Dermatitis caused by neck threadworm microfilariae, *Onchocerca* sp.

**Precautions:** All horses should be included in a regular parasite control program with particular attention being paid to mares, foals, and yearlings. Foals should be treated initially at 6 to 8 weeks of age, and routine treatment repeated as appropriate. Consult your veterinarian for a control program to meet your specific needs. Ivermax® (ivermectin) Equine Paste Dewormer effectively controls gastrointestinal nematodes and bots of horses. Regular treatment will reduce the chances of verminous arteritis caused by *Strongylus vulgaris*.

**Product Advantages:** Broad-spectrum Control – Ivermax® (ivermectin) Equine Paste Dewormer kills important internal parasites, including bots and the arterial stages of *S. vulgaris*, with a single dose. Ivermax® Equine Paste Dewormer is a potent antiparasitic agent that is neither a benzimidazole nor an organophosphate.

**Animal Safety:** Ivermax® (ivermectin) Equine Paste Dewormer may be used in horses of all ages, including mares at any stage of pregnancy. Stallions may be treated without adversely affecting their fertility.